

Simple quantitative estimate of the derivative discontinuity and the energy gap in atoms and molecules

F. P. Rosselli,¹ A. B. F. da Silva,² and K. Capelle^{3,*}

¹*Divisão de Metrologia de Materiais, Instituto Nacional de Metrologia, Normalização e Qualidade Industrial (Inmetro), Rio de Janeiro 25245-020, Brazil*

²*Departamento de Química e Física Molecular, Instituto de Química de São Carlos, Universidade de São Paulo, CP 780, São Carlos, 13560-970 SP, Brazil*

³*Departamento de Física e Informática, Instituto de Física de São Carlos, Universidade de São Paulo, CP 369, 13560-970 São Carlos, SP, Brazil*

(Dated: February 2, 2008)

The derivative discontinuity of the exchange-correlation functional of density-functional theory is cast as the difference of two types of electron affinities. We show that standard Kohn-Sham calculations can be used to calculate both affinities, and that their difference benefits from substantial and systematic error cancellations, permitting reliable estimates of the derivative discontinuity. Numerical calculations for atoms and molecules show that the discontinuity is quite large (typically several eV), and significantly and consistently improves the agreement of the calculated fundamental energy gap with the experimental gaps. The percentage error of the single-particle gap is reduced in favorable cases by more than one order of magnitude, and even in unfavorable cases by about a factor of two.

PACS numbers: 31.15.E-, 31.15.es, 31.15.-p, 71.15.Mb

From spectroscopy to transport, there is hardly any property of a quantum many-particle system that does not in some way depend on whether there is a gap in the energy spectrum, and what the size of this gap is. The fundamental gap is a ground-state property of the N -body system, defined [1, 2, 3] in terms of the ground-state energy $E(N)$ as $E_g = [E(N-1) - E(N)] - [E(N) - E(N+1)]$, where $E(N-1) - E(N)$ is the energy change upon removing the N 'th particle from the N particle system and $E(N) - E(N+1)$ that upon removing the $N+1$ st particle from the $N+1$ particle system. In a noninteracting system, this definition reduces to the familiar energy gap between single-particle levels.

For interacting systems, approximate many-body calculations of total energies and energy gaps are typically performed within the framework of density-functional theory (DFT) [1, 2, 3] or, for small finite systems, Hartree-Fock (HF) and post-Hartree-Fock methods. DFT provides, in addition to the ground-state density and related quantities, also a set of single-particle eigenvalues, the so-called Kohn-Sham (KS) spectrum. The difference between the energy of the highest occupied and lowest unoccupied of these single-particle levels is the KS gap, which in extended systems become the band structure gap. Neither the KS nor the HF single-particle gaps correspond to the experimental gap, the former typically being too small and the latter too large.

Generally, one can write $E_g = E_g^{KS} + \Delta_{xc}$, which defines Δ_{xc} as the difference between the exact fundamental gap and the exact KS single-particle gap. In atomic physics and quantum chemistry, the importance of a

nonzero Δ_{xc} for chemical hardness is well known [2, 4, 5]. Neglect of Δ_{xc} has also been shown to lead to large errors in the calculation of Rydberg excitations [6, 7], charge-transfer processes and transport [8, 9, 10, 11], and the ionization probability of atoms exposed to strong electromagnetic fields [12]. In semiconductors, approximate energy gaps calculated in DFT often drastically underestimate the experimental gap [13, 14]. In Mott insulators, in particular, the entire gap is due to Δ_{xc} [15, 16]. Recently it was pointed out that a similar discontinuity problem can also appear in the spin-DFT calculation of spin gaps in materials relevant for spintronics [17]. The question whether the neglect of Δ_{xc} or the error in $E_g^{KS,approx}$ is responsible for the underestimate of the band gap in solids is considered in a standard textbook in the field to be ‘*of tremendous practical importance*’ [1], and the calculation of Δ_{xc} is ranked in a recent monograph as ‘*certainly one of the major outstanding problems in contemporary DFT*’ [3], but no general answer is known.

In the present paper we draw attention to an alternative representation of Δ_{xc} , which casts it as a difference of single-particle eigenvalues, similar to the KS gap. We point out that this relation provides a simple physical interpretation of the elusive xc discontinuity, and use it to estimate xc discontinuities of atoms and molecules. The resulting correction Δ_{xc}^{est} substantially improves agreement with experimental fundamental gaps, reducing the percentage deviation from experiment by more than a factor of 10 in favorable cases and by about a factor of two in less favorable ones.

The standard representation of Δ_{xc} is based on ensemble DFT for open systems, where all three quantities in $E_g = E_g^{KS} + \Delta_{xc}$ can be related to derivative discontinuities of universal density functionals [15, 18, 19]. The fundamental gap is the derivative discontinuity of the to-

*Electronic address: capelle@ifsc.usp.br

TABLE I: Experimental gap, $E_g^{exp} = I^{exp} - A^{exp}$, KS single-particle gap, E_g^{KS} , and KS gap corrected by adding our estimate for the discontinuity, $E_g^{DFT} = E_g^{KS} + \Delta_{xc}^{est}$, for atoms *Li* ($Z = 3$) to *Ca* ($Z = 20$), with exception of the noble gases. The values for Δ_{xc}^{est} were obtained from Eq. (7) using the B88LYP functional and the 6-311G(d) basis sets. All data in eV.

Z	3	4	5	6	7	8	9	11	12	13	14	15	16	17	19	20	rms
	Li	Be	B	C	N	O	F	Na	Mg	Al	Si	P	S	Cl	K	Ca	error
E_g^{KS}	1.40	3.56	0.614	0.598	3.76	0.860	0.820	0.917	3.38	0.272	0.188	2.01	0.215	0.146	0.601	2.36	7.67
E_g^{DFT}	4.59	9.01	8.12	10.1	14.4	13.2	15.1	4.42	7.19	5.09	6.43	8.45	7.64	8.88	3.63	5.32	0.606
E_g^{exp}	4.77	9.32	8.02	10.0	14.5	12.2	14.0	4.59	7.65	5.55	6.76	9.74	8.28	9.36	3.84	6.09	-

tal energy

$$E_g = \left. \frac{\delta E[n]}{\delta n(\mathbf{r})} \right|_{N+\eta} - \left. \frac{\delta E[n]}{\delta n(\mathbf{r})} \right|_{N-\eta}, \quad (1)$$

the KS single-particle gap that of the noninteracting kinetic energy

$$E_g^{KS} = \left. \frac{\delta T_s[n]}{\delta n(\mathbf{r})} \right|_{N+\eta} - \left. \frac{\delta T_s[n]}{\delta n(\mathbf{r})} \right|_{N-\eta}, \quad (2)$$

and the remaining piece, Δ_{xc} , that of the xc energy

$$\Delta_{xc} = \left. \frac{\delta E_{xc}[n]}{\delta n(\mathbf{r})} \right|_{N+\eta} - \left. \frac{\delta E_{xc}[n]}{\delta n(\mathbf{r})} \right|_{N-\eta} = v_{xc}^+(\mathbf{r}) - v_{xc}^-(\mathbf{r}). \quad (3)$$

In these equations η stands for an infinitesimal variation of the system's particle number. Equation (3) shows that Δ_{xc} is a system-dependent shift of the xc potential $v_{xc}(\mathbf{r})$ as it passes from the electron-poor to the electron-rich side of integer N .

These three relations are useless to calculate gaps from most currently available approximate density functionals, which typically have no discontinuities. For two of the three quantities above, alternative ways of calculation, more useful in practice, are widely known. Total energies are calculated easily from DFT, so that one can employ the definition of ionization energy $I = E(N-1) - E(N)$ and electron affinity $A = E(N) - E(N+1)$ to calculate the fundamental gap from

$$E_g = E(N+1) + E(N-1) - 2E(N) = I - A. \quad (4)$$

Single-particle energies are obtained from the KS equation, as a byproduct of calculating the total energies, and yield the KS single-particle gap

$$E_g^{KS} = \epsilon_{N+1}(N) - \epsilon_N(N), \quad (5)$$

where $\epsilon_M(N)$ denotes the M 'th eigenvalue of the N -electron system. The third term, the xc discontinuity Δ_{xc} , has resisted all attempts of describing it directly by common density functionals, such as LDA and GGA, which are continuous as a function of N and thus have no xc discontinuity.

However, we note that I and A can be calculated in DFT not only from ground-state energies, but also from

single-particle eigenvalues, via $I = -\epsilon_N(N)$ and $A = -\epsilon_{N+1}(N+1)$ (the analogue of Koopmans' theorem in DFT) [1, 20]. By using these relations and the definition of E_g^{KS} one finds, upon combining Eqs. (4) and (5) [21],

$$E_g = \epsilon_{N+1}(N) - \epsilon_N(N) + \Delta_{xc} = \epsilon_{N+1}(N+1) - \epsilon_N(N), \quad (6)$$

which implies

$$\Delta_{xc} = \epsilon_{N+1}(N+1) - \epsilon_{N+1}(N) = A_{KS} - A. \quad (7)$$

In the last step we used the fact that the affinity of the KS system, A_{KS} , is simply the negative of the energy of the lowest unoccupied orbital.

One way to estimate Δ_{xc} is by subtracting Eq. (5) from (4). In the following, we use Eq. (7), which is equivalent in principle, but simpler in practice [22]. It also provides an intuitive interpretation of the discontinuity: in an interacting system, the electrons repel, hence the energy cost of removing the outermost electron from the negative species (which is measured by the electron affinity) is reduced, $A < A_{KS}$, and a positive Δ_{xc} results.

If the right-hand side of Eq. (7) could be calculated exactly, this procedure would determine the exact xc discontinuity. An estimate of Δ_{xc} is thus obtained by using in (7) the KS eigenvalues obtained in two approximate KS calculations, one for the neutral species, the other for the anion.

Such approximate calculations involve two distinct types of errors, one associated with the approximations used for the xc functional, the other with the finite size of the basis set. As a consequence, each of the two affinities in Eq. (7) is predicted wrongly by standard combinations of functionals and basis sets. Typically, the self-interaction error inherent in common LDA and GGA type functionals shifts the eigenvalues up, in some cases so much that the anions become unbound [23]. On the other hand, the finiteness of the basis set artificially stabilizes the anion [23]. As a consequence of this error cancellation, practical methods for calculating affinities from LDA and GGA are available [23, 24, 25, 26].

Our key argument is that derivative discontinuities are protected from functional errors and basis-set errors by a distinct additional error compensation, independent of the one just described. Namely, Eq. (7) casts Δ_{xc} as a difference of two affinities. Even if each is predicted wrongly on its own, their difference is expected to benefit from substantial additional error cancellation. In fact,

if all KS eigenvalues are shifted up by roughly the same amount, energy differences are preserved, and even positive eigenvalues (unbound anions) can provide reasonable discontinuities for the bound neutral system.

We call this calculation of Δ_{xc} by means of Eq. (7) an estimate, and not an approximation, to stress that it exploits an error cancellation that is hard to quantify *a priori*. However, in recent work on models of harmonically confined systems [27] this estimate was found to lead to significantly improved gaps. Here we explore the performance of Eq. (7) in *ab initio* calculations for atoms and molecules.

In Table I we compare, for 16 light atoms, the experimental gap, the KS single-particle gap, and the DFT gap, defined as the sum of the KS gap and the estimated xc discontinuity. The KS calculations were performed with the GAUSSIAN 98 [28] program, using the B88-LYP functional and the 6-311G(*d,p*) basis sets. Table I shows that the error of the KS gap is significantly and consistently reduced by adding the estimated xc discontinuity to the KS gap, dropping by more than an order of magnitude — from 7.67eV to 0.606eV — over the data set in Table I. This large drop, together with the fact that the improvement is systematic (obtained not only on average, but in every individual case), strongly suggests that Eq. (7) is a reliable and useful way of obtaining the discontinuity.

Figure 1 is a plot of the data in Table I, revealing that Δ_{xc} roughly follows the atomic shell structure. Particularly small discontinuities are found for atoms with one electron outside a closed shell, such as *Li* and *Na*. The largest discontinuities are, however, not observed for closed-shell systems but for systems one electron short of a closed shell, as is seen comparing *F* with *Ne* or *Cl* with *Ar*. We interpret this by means of Eq. (7) as a consequence of the fact that Δ_{xc} is related to two affinities, which involve negative species with one additional electron, leading to a closed shell for *F*[−] and *Cl*[−].

Specifically for the *Be* atom, we can further compare with independent theoretical expectations, because the discontinuity of this atom has previously been estimated by Jones and Gunnarsson (JG) [29] by comparing the experimental gap to a near-exact KS gap obtained earlier by Pedroza and Almbladh [30] from CI densities and approximate inversion of the KS equation. Our value $\Delta_{xc}^{Be} = 5.5\text{eV}$ is encouragingly close to $\Delta_{xc}^{Be,JG} = 5.7\text{eV}$.

Next, we turn to molecules. In Table II we compare our estimate of Δ_{xc} to many-body values of Δ_{xc} and to experimental fundamental gaps. The many-body discontinuity is obtained [31] by performing coupled-cluster calculations to generate a near-exact density, followed by inversion of the KS equation to obtain the corresponding near-exact KS potential, solution of the KS equation with that potential to obtain the near-exact KS gap, and subtraction of that gap from the experimental fundamental gap [31]. The first step is impractical for larger systems, whereas the last step involves using the experimental gap, which makes the method empirical. For these




FIG. 1: Plot of the data in Table I: Kohn-Sham single-particle gap (triangles), experimental fundamental gap (open squares) and Kohn-Sham single-particle gap corrected by adding our estimate of the discontinuity (full circles), for 16 light atoms. The lines are guides for the eye, illustrating that Δ_{xc}^{est} recovers not only the overall value, but also fine details of the behaviour as a function of *Z* that were lost in the KS gap.

reasons, the simple estimate obtained from Eq. (7) may constitute a useful alternative, provided it turns out to be sufficiently reliable.

In fact, the estimated value of Δ_{xc}^{est} depends on the chosen basis set and functional, as well as on whether the anion geometry is separately optimized (leading to adiabatic affinities and discontinuities) or held fixed at that of the neutral species (vertical affinities and discontinuities). Tests of different combinations of methodologies indicate that best (*i.e.*, closest to experiment) gaps are obtained if the discontinuity is calculated from vertical affinities.

Table II shows that the systems fall in two classes. For one class, comprising *CO*, *H₂CO*, *H₂S*, *HCN*, *N₂*, *PH₃*, *Cl₂*, *SO₂*, and *C₂H₄*, the gap error with respect to experiment is reduced by a similar margin as for atoms, or even more, and estimated and calculated Δ_{xc} agree well. For the other class, comprising *H₂O*, *NH₃*, *HF*, *CH₄* and perhaps *C₂H₂*, the percentage error of the gap drops by a factor of two, instead of by one order of magnitude, and the estimate recovers about 50% of the many-body value of Δ_{xc} . An empirical indicator of which class a system belongs to is the sign of the KS affinity: if this is negative (*i.e.*, the lowest unoccupied KS orbital has positive eigenvalue) the system belongs to class II. We note that this is not a stability criterium because it employs the (unphysical) KS affinity and because for all molecules in Table II (with exception of *Cl₂* and *SO₂*) the anionic species is experimentally unstable. Rather, it indicates a partial loss of the error cancellation on which our use of Eq.(7) is based. Nevertheless, even in these "unfavorable" cases, the estimate still provides a systematic improvement on the KS gap, reducing the error with respect to experiment by a factor of two. For both classes, Δ_{xc}^{est} clearly

TABLE II: Comparison of calculated and experimental gaps for small molecules. First column: experimental gap. Second column: KS single-particle gap. Third column: percentage deviation of KS gap from experimental gap. Fourth column: derivative discontinuity estimated from single-particle eigenvalues, obtained from the B88-LYP functional on the 6-311G(d,p) basis sets. Fifth column: DFT gap. Sixth column: percentage deviation of DFT gap from experimental gap. Seventh column: many-body estimate of the derivative discontinuity. All values in eV.

system	$E_g^{expt.}$, Ref. [32]	E_g^{KS}	% dev.	Δ_{xc}^{est} , Eq.(7)	$E_g^{DFT} = E_g^{KS} + \Delta_{xc}^{est}$	% dev.	Δ_{xc} , Ref. [31]
CO	15.8	7.05	-55.4	9.04	16.09	1.83	8.44
H ₂ CO	12.4	3.66	-70.5	8.31	11.97	-3.43	8.16
H ₂ S	12.6	5.68	-54.9	5.33	11.01	-12.6	6.53
HCN	15.9	8.06	-49.3	8.24	16.30	2.53	7.89
N ₂	17.8	8.24	-53.7	9.74	17.98	1.15	9.25
PH ₃	11.9	6.51	-45.3	5.25	11.76	-1.18	5.99
Cl ₂	9.20	2.38	-74.1	7.24	9.620	4.57	-
SO ₂	11.2	3.25	-71.0	7.94	11.19	-0.09	-
C ₂ H ₄	12.3	5.76	-53.2	7.10	12.86	4.57	6.53
C ₂ H ₂	14.0	6.97	-50.2	5.34	12.32	-12.0	7.08
H ₂ O	19.0	6.47	-66.0	6.15	12.61	-33.6	11.4
NH ₃	16.3	6.01	-63.1	5.56	11.58	-29.0	10.1
HF	22.0	8.63	-60.8	6.82	15.45	-29.8	11.7
CH ₄	20.5	10.3	-49.9	4.88	15.15	-26.1	11.4

provides a quantitative correction to single-particle gaps, which may be useful in improving, e.g., the DFT description of chemical hardness [4, 5] or of transport through single molecules [8, 9].

In summary, we have cast the derivative discontinuity of DFT as a difference of two affinities, Eq. (7), and used approximate KS calculations of these to obtain estimates

for the discontinuity in atoms and molecules. Our results are consistent with previous results, where available, and significantly and consistently reduce the error between calculated and measured fundamental gaps.

This work was supported by FAPESP, CAPES and CNPq.

-
- [1] R. M. Dreizler and E. K. U. Gross, *Density Functional Theory* (Springer-Verlag, Berlin, 1990).
- [2] R. G. Parr and W. Yang, *Density-Functional Theory of Atoms and Molecules* (Oxford University Press, Oxford, 1989).
- [3] G. F. Giuliani and G. Vignale, *Quantum Theory of the Electron Liquid* (Cambridge University Press, 2005).
- [4] A. Görling and M. Levy, Phys. Rev. A **52**, 4493 (1995).
- [5] D. J. Tozer and F. De Proft, J. Phys. Chem. A **109**, 8923 (2005).
- [6] A. Wasserman, N. T. Maitra and K. Burke, Phys. Rev. Lett. **91**, 263001 (2003).
- [7] D. J. Tozer and N. C. Handy, J. Chem. Phys. **109**, 10180 (1998).
- [8] C. Toher, A. Filippetti, S. Sanvito and K. Burke, Phys. Rev. Lett. **95**, 146402 (2005).
- [9] S.-H. Ke, H. U. Baranger and W. Yang, J. Chem. Phys. **126**, 201102 (2007).
- [10] D. J. Tozer, J. Chem. Phys. **119**, 12697 (2003).
- [11] Q. Wu and T. Van Voorhis, Phys. Rev. A **72**, 024502 (2005).
- [12] M. Lein and S. Kümmel, Phys. Rev. Lett. **94**, 143003 (2005).
- [13] R. W. Godby, M. Schlüter and L. J. Sham, Phys. Rev. Lett. **56**, 2415 (1986).
- [14] M. Grüning, A. Marini and A. Rubio, J. Chem. Phys. **124**, 154108 (2006).
- [15] J. P. Perdew and M. Levy, Phys. Rev. Lett. **51**, 1884 (1983).
- [16] N. A. Lima, L. N. Oliveira, and K. Capelle, Europhys. Lett. **60**, 601 (2002).
- [17] K. Capelle and G. Vignale, Phys. Rev. Lett. **86**, 5546 (2001).
- [18] L. J. Sham and M. Schlüter, Phys. Rev. Lett. **51**, 1888 (1983).
- [19] W. Kohn, Phys. Rev. B **33**, 4331 (1986).
- [20] Q. Wu, P. W. Ayers, and W. Yang, J. Chem. Phys. **119**, 2978 (2003).
- [21] L. J. Sham and M. Schlüter, Phys. Rev. B **32**, 3883 (1985).
- [22] Eq. (7) does not require calculations for the $N-1$ particle system, and also avoids numerically troublesome calculation of Δ_{xc} as difference of much larger numbers.
- [23] N. Rösch and S. B. Trickey, J. Chem. Phys. **106**, 8940 (1997).
- [24] J. M. Galbraith and H. F. Schaefer, III, J. Chem. Phys. **105**, 862 (1996).
- [25] G. de Oliveira, J. M. L. Martin, F. de Proft, and P. Geerlings, Phys. Rev. A **60**, 1034 (1999).
- [26] O. A. Vydrov and G. E. Scuseria, J. Chem. Phys. **122**,

- 184107 (2005).
- [27] K. Capelle, M. Borgh, K. Karkkainen and S. M. Reimann, Phys. Rev. Lett. **99**, 010402 (2007).
- [28] *Gaussian 98*, M. J. Frisch et al., Gaussian Inc., Pittsburgh, PA (1998).
- [29] R. O. Jones and O. Gunnarsson, Rev. Mod. Phys. **61**, 689 (1989).
- [30] C.-O. Almbladh and A. C. Pedroza, Phys. Rev. A **29**, 2322 (1984).
- [31] M. J. Allen and D. J. Tozer, Mol. Phys. **100**, 433 (2002).
- [32] R. G. Pearson, Inorg. Chem. **27**, 734 (1988).